

ST102 Week 7

Def. (Moments of a random variable)

For $k = 1, 2, \dots$ define

i) the k th moment about zero as $\mu_k := E(\bar{X}^k)$

ii) the k th central moment as $\mu'_k := E[(\bar{X} - E(\bar{X}))^k]$

Remark. Clearly, $\mu_1 = E(\bar{X})$; $\mu'_2 = \text{Var}(\bar{X})$

Def. (Moment generating function / mgf)

The mgf of a discrete r.v. \bar{X} is defined as:

$$M_{\bar{X}}(t) = E(e^{t\bar{X}}) = \sum_{x \in \mathbb{D}} e^{tx} p(x)$$

Remark. $M_{\bar{X}}(t)$ is a function in t , with no randomness

Prop. (Useful properties of a mgf)

i) $M_{\bar{X}}'(0) = E(\bar{X})$ & $M_{\bar{X}}''(0) = E(\bar{X}^2)$

... & $M_{\bar{X}}^{(k)}(0) = E(\bar{X}^k)$, $k \geq 1$

ii) $\text{Var}(\bar{X}) = E(\bar{X}^2) - (E\bar{X})^2 = M_{\bar{X}}''(0) - [M_{\bar{X}}'(0)]^2$

iii) $M_{\bar{X}}(t) = M_Y(t) \Rightarrow \bar{X} \& Y \text{ share the same distribution}$

iv) $M_{\bar{X}+b}(t) = e^{bt} M_{\bar{X}}(at)$

v) For independent r.v.'s $\bar{X}_1, \dots, \bar{X}_n$,

$$M_{\sum_{i=1}^n \bar{X}_i}(t) = \prod_{i=1}^n M_{\bar{X}_i}(t)$$

Def. i) Continuous r.v.

ii) probability density function $f(x)$ (p.d.f.)

R.K. $P(\underline{X} = x_0) = 0, \forall x_0 \in \mathbb{Q}$

$$\text{iii) } P(a < x < b) = \int_a^b f(x) dx$$

iv) Necessary conditions

$$\begin{cases} f \geq 0 & \forall x \\ \int_{-\infty}^{+\infty} f(x) dx = 1 \end{cases}$$

v) cumulative distribution function (c.d.f.)

$$F(x) := P(\underline{X} \leq x) = \int_{-\infty}^x f(t) dt$$

$$\text{R.K. } f(x) = \frac{d}{dx} F(x);$$

$$P(a < x < b) = \int_a^b f(x) dx = F(b) - F(a)$$

$$\text{vi) } E(X) = \int_{-\infty}^{+\infty} x f(x) dx$$

$$E(g(X)) = \int_{-\infty}^{+\infty} g(x) f(x) dx$$

$$\text{Var}(X) = E[(X - E(X))^2] = \int_{-\infty}^{+\infty} (x - E(x))^2 f(x) dx$$

$$= E(X^2) - [E(X)]^2$$

$$sd(X) = \sqrt{\text{Var}(X)}$$